arXiv:physics/0703258v2 [physics.optics] 8 May 2010

On the physical nature of the optical precursor phenomenon

V.A. Kuz'menko[1]

Troitsk Institute for Innovation and Fusion Research,
Troitsk, Moscow region, 142190, Russian Federation.

Abstract

Physical phenomenon of the optical precursor is a beautiful manifestation of strong inequality of forward and reversed transitions in optics.

PACS number: 42.50.Md, 42.50.Ct

Present day quantum physics is overloaded by inferior mathematical descriptions. However, too frequently such mathematical models can not help and only replace understanding of a physical nature by the description of the studied phenomena.

The striking example of such situation is the phenomenon of so-called optical precursor. Its mathematical description for the first time was proposed nearly one hundred years ago [1], but high quality experimental results were obtained only recently [2 - 4]. The used mathematical model allows exactly to describe the shape of the precursor, but it is useless for understanding the nature of observed phenomenon. It is supposed in [4] that: "an optical precursor is the propagation of the leading edge of the step pulse that travels at the speed of light in vacuum because the atoms have finite response time - they cannot respond to the leading edge immediately". So, it is assumed that the time of a photon absorption event is insufficiently short. Of course, this is an absurd supposition because of the characteristic time of the observed optical precursor is near 10 nc. However, the physicists are in a hopeless position: any other possible physical explanation looks like even more absurdly than the present one.

But the physical nature of the observed optical precursor phenomenon (as also of many other phenomena in optics) is easily explained on the base of the concept of inequality of forward and reversed processes in optics [5]. According to this concept the differential cross-sections of forward process of photon absorption and following reversed process of a stimulated emission of the photon by the same atom are not equal. Figure 1 shows the supposed shapes of corresponding dependences of cross-sections from different physical parameters.

The physical nature of the optical precursor phenomenon becomes clear from this figure. At the first stage a photon absorption process takes place. It has some probability or cross-section. At the same time the reversed process of stimulated emission of the photon proceeds. It has much greater differential cross-section. As a result, the total observed absorption of photons is small. Stimulated emission is the same statistical process. If it does not take place

---

[1] Electronic address: kuzmenko@triniti.ru

immediately, the different dephasing (in wide sense) and states mixing processes take place. As a result, the differential cross-section of the backward stimulated emission process strongly drops, the atoms "stick" in the excited state and the observed total absorption of photons quickly grows. At the end of the pulse due to the rephrasing processes some excess of stored in the gas medium energy is emitted without any inversion of the state's populations.

It may seems that the concept of inequality of forward and reversed processes contradicts to well known phenomenon of Rabi oscillations of population of states in two-level quantum system under action of resonance radiation, which looks like as a demonstration of full symmetry of forward and backward processes. However, such contradiction is absent. For Rabi oscillations it is important the equality of total (integral) cross-sections of forward and backward processes (this equality is not disputed). In contrast, for optical precursor phenomenon and for the so-called adiabatic population transfer phenomenon [6] (during the sweeping of resonance conditions) it is more important the inequality of differential cross-sections of forward and reversed transitions.

The task of full explanation and more correct mathematical description of the optical precursor phenomenon demands extensive information about differential cross-sections of forward, backward, reversed transitions and its dependence from a number of physical parameters. Now such information is absent. Present day experimental results [2 - 4] have mainly demonstrational character. We need now in experimental study of relative intensity and time length of the optical precursor from the intensity of laser radiation, gas temperature and so on in as possible wide range of experimental parameters. It is just probably that the extensive experimental study of the optical precursor phenomenon will give in a future most exact quantitative information about differential cross-sections of a photon interaction with atoms and molecules.

References

[1] L. Brillouin, *"Wave Propagation and Group Velocity"*, Academic Press, New York, (1960).

[2] H. Jeong, A.M.C. Dawes, and D.J. Gauthier, Phys. Rev. Lett. **96**, 143901 (2006).

[3] D. Wei, J.F. Chen, M.M.T. Loy, G.K.L. Wong, and S. Du, Phys. Rev. Lett. **103**, 093602 (2009).

[4] J.F. Chen, S. Wang, D. Wei, M.M.T. Loy, G.K.L. Wong, and S. Du, Phys. Rev. A **81**, 033844 (2010).

[5] V.A. Kuz'menko, e-print, physics/0306148.

[6] L. Allen, J.H. Eberly, *"Optical Resonance and Two-Level Atoms"*, John Wiley and Sons, (1975).



Figure1 The supposed shapes of dependences of cross-sections for forward (1), backward and reversed (2) transitions from such parameters as the frequency (ν), phase, angel (θ), position of the particle (x).